arXiv:1207.2132v1 [math.MG] 9 Jul 2012

Mapping Class Groups have finite Assouad-Nagata dimension

David Hume*

July 10, 2012

Abstract

We improve a result of Bestvina, Bromberg and Fujiwara to prove that mapping class groups quasi-isometrically embed into finite products of tree-graded spaces whose pieces are curve complexes. From this we deduce that the Assouad-Nagata dimension of these groups is finite and provide explicit embeddings into ℓ^p spaces with compression exponent 1.

1 Introduction

The notion of asymptotic dimension was introduced by Gromov [Gro93] as a way of extending the notion of dimension to infinite groups. Assouad-Nagata dimension was formulated by Assouad (under the name Nagata dimension) [Ass82] and in more recent work it has been seen as a stronger linearly-controlled version of asymptotic dimension. There are natural examples of finitely generated groups with finite asymptotic dimension but infinite Assouad-Nagata dimension [BDL].

Finite Assouad-Nagata dimension implies many useful properties for a group

- bounds the topological dimension of asymptotic cones [DH08],
- provides certain Lipschitz extension properties [BDHM09, LS05]
- guarantees ℓ^p compression exponent 1 for all $p \geq 1$ [Gal08].

Very few classes of finitely generated groups are known to have finite Assouad-Nagata dimension: abelian groups, hyperbolic groups and certain wreath products do [Now07, BDL], also bounds can be found for groups G obtained from short exact sequences $1 \to N \to G \to K \to 1$ whenever $(N, d_G|_N)$ and K have finite Assouad-Nagata dimension [BDLM08]. It is not known whether all nilpotent groups have finite Assouad-Nagata dimension, though it certainly does not hold for metabelian groups, for example the wreath product $\mathbb{Z} \wr \mathbb{Z}$ contains quasi-isometrically embedded copies of $\mathbb{Z}^n$ for each n.

Having finite Assouad-Nagata dimension is equivalent to admitting a quasisymmetric embedding into a product of trees [Sch05], but it is unknown whether this

---

[*] University of Oxford, Mathematical Institute, 24-29 St Giles, Oxford OX1 3LB, United Kingdom. Email: hume@maths.ox.ac.uk

is strictly weaker than having ℓ^p compression exponent 1 for all $p \in [1, \infty)$. We recall that the ℓ^p compression exponent of a metric space (X, d) is the supremum over $\alpha \in [0, 1]$ such that there exists an embedding ϕ of X with

$$d(x,y)^\alpha \preceq \|\phi(x) - \phi(y)\|_p \preceq d(x,y),$$

where $f \preceq g$ means there exists some constant C such that $f(n) \leq Cg(n) + C$ for all n. This is known to hold for a slightly larger class of groups than just those currently known to have finite Assouad-Nagata dimension: polycylic groups, free products of groups with compression exponent 1 and certain wreath products but not others [Tes11, Hum11, NP11]. A slightly weaker statement is true for all relatively hyperbolic groups with parabolic subgroups of compression exponent 1, they are known to have ℓ^p compression exponent 1 for all $p > 1$ [Hum11]. Even in cases where the compression exponent is known it is difficult in general to actually provide an embedding exhibiting this.

One of the most interesting classes of finitely generated groups are mapping class groups - homeomorphisms of surfaces considered up to isotopy - due to their close connections with geometry, topology and group theory and their similarities with $Out(F_n)$ and lattices in higher rank semisimple Lie groups. In light of our results it is an interesting question to ask whether these groups necessarily have compression exponent 1 or even finite Assouad-Nagata dimension.

In the class of hyperbolic spaces, the quasi-isometry class containing all trees provides an important subclass defined by the bottleneck property [Man05], similarly, within relatively hyperbolic spaces the quasi-isometry class containing all tree-graded spaces (the metric formulation of a free product of groups) is a subclass with additional structural properties, most notably in our setting the passing of finite (asymptotic) Assouad-Nagata dimension and ℓ^p compression 1 from pieces to the whole space [BH, Hum11].

Bestvina, Bromberg and Fujiwara prove that a certain list of axioms allow the construction of quasi-trees on which many groups (mapping class groups included) act. In particular, they obtain a list of finitely many quasi-trees with vertices in $1-1$ correspondence with subsurfaces of a particular type, such that the direct product of these quasi-trees “blown up” at each vertex by a curve complex $\mathcal{C}(W)$ contains a quasi-isometric copy of the mapping class group [BBF].

In this paper, we prove that each of the factors in this product is quasi-isometric to a tree-graded space, and indeed each hyperbolic “blown-up” quasi-tree satisfying the Bestvina-Bromberg-Fujiwara axioms with one-ended pieces is a quasi tree-graded space. More precisely,

Theorem 1. (4.1) *Consider a quasi-tree of spaces $\mathcal{C}(\mathbf{Y})$ satisfying the axioms listed in* [BBF]*. If $\mathcal{C}(\mathbf{Y})$ is hyperbolic and all pieces $\mathcal{C}(Y)$ are one-ended then $\mathcal{C}(\mathbf{Y})$ is quasi-isometric to a tree-graded space $\mathcal{T}(\mathbf{Y})$ with set of pieces $\{\mathcal{C}(Y) \mid Y \in \mathbf{Y}\}$.*

This result increases interest in determining which curve complexes are one-ended. Indeed if all curve complexes were quasi-trees then this result would be redundant by [BBF], but this is far from the case [Sch11]. The hyperbolicity of $\mathcal{C}(\mathbf{Y})$ in the case of the mapping class group embedding comes from the Masur-Minksy theorem that curve complexes are δ-hyperbolic [MM99, BBF].

From this we obtain an explicit quasi-isometric embedding of each mapping class group into a product of finitely many tree-graded spaces each of which has curve complexes of homeomorphic subsurfaces of S as its pieces,

$$MCG(S) \to \prod \mathcal{C}(\mathbf{Y}) \to \prod \mathcal{T}(\mathbf{Y}).$$

We then obtain the main theorem.

Theorem 2. (4.5) *The mapping class group of any compact surface has finite Assouad-Nagata dimension.*

This was previously only known in low complexity cases, where the mapping class group is virtually free [Beh04].

In addition to theorem 1, for the proof of theorem 2 we also require the theorem of Bell-Fujiwara which states the curve complexes have finite asymptotic dimension, but, in fact, their exact argument proves the stronger result of finite Assouad-Nagata dimension [BF08].

Previously there was little information concerning how mapping class groups may embed into Banach spaces. Finite asymptotic dimension does imply coarse embeddability into Hilbert spaces, so mapping class groups satisfy the strong Novikov and Coarse Baum-Connes conjectures - the Novikov conjecture had already been granted independently by work of Hamenstädt, Kida and Behrstock-Minsky. Kida, moreover, proves that mapping class groups are exact and hence have Yu's property (A), which also follows from finite asymptotic dimension. [BBF, HR00, Yu00, Ham09, Kid08, BM11]

Theorem 2 implies that $MCG(S)$ has ℓ^p compression exponent 1 for all p [Gal08]. As a corollary of our methods we provide explicit embeddings of mapping class groups into ℓ^p spaces exhibiting compression exponent 1 (4.10), by first proving the result for curve complexes (4.6) using Bowditch's results on the Masur-Minksy definition of tight geodesics [Bow08, MM00]. This provides the first examples of hyperbolic graphs without bounded growth at any scale (other than quasi-trees) known to have compression exponent 1. We then extend this embedding to tree-graded spaces with curve complexes as pieces. Both steps use previous work of the author [Hum11].

As an outline of the paper, in section 2 we recall the necessary preliminaries required from [BBF], then define the appropriate tree-graded spaces $\mathcal{T}(\mathbf{Y})$ in section 3. Section 4 contains the proofs of Theorems 1 and 2 as well as details of explicit embeddings into ℓ^p spaces.

2 Properties of quasi-trees of metric spaces

We first recall the features of the quasi-trees of metric spaces defined in [BBF] that are required to progress.

- $\mathcal{C}(\mathbf{Y})$ is a metric graph in which the pieces $\mathcal{C}(Y)$ are totally geodesically embedded simplicial graphs. (We will assume throughout that the spaces $\mathcal{C}(Y)$ are all one-ended.)
- There are projections π_Y which map any $X \in \mathbf{Y} \setminus \{Y\}$ to a subset of $\mathcal{C}(Y)$ with diameter bounded by L.

- There exist standard paths between any two pieces, the internal pieces of which are written as an ordered set $Y_K(X,Z)$ where two spaces $\mathcal{C}(X)$ and $\mathcal{C}(Z)$ are joined by a collection of edges of length L with endpoints in $\pi_Y(X) \times \pi_X(Y)$ if and only if $Y_K(X,Z) = \emptyset$.

Moreover, we have the following collection of bottlenecks.

Lemma 2.1. *Let $X, Z \in \mathbf{Y}$ with $X \neq Z$ and let $Y \in Y_K(X,Z) \cup \{Z\}$. There is some point $w_Y \in \pi_Y(X)$ such that all paths from $\mathcal{C}(X)$ to $\mathcal{C}(Z)$ pass within distance $9L$ of w_Y.*

Proof: For $Y \in Y_K(X,Z)$ the result follows directly from [BBF, Lemma 3.9] with $w_Y \in \pi_Y(X)$. We deal only with the case $Y = X$, as the case $Y = Z$ is obtained by reversing the orientation of paths.

Suppose first that $Y_K(X,Z) = \emptyset$. As the space $\mathcal{C}(Z)$ is one-ended we may choose a point $z \in \mathcal{C}(Z)$ such that $Y_K(X,z) = \{Z\}$, using the axiom $\operatorname{diam}(\pi_Z(X)) \leq K$. Take any path P from some $x \in \mathcal{C}(X)$ to z. By [BBF, Lemma 3.9] there is a point w lying on P such that $d_{\mathcal{C}(\mathbf{Y})}(w, \pi_Z(X)) \leq 7L$. As any $\mathcal{C}(X)$ to $\mathcal{C}(Z)$ path not meeting $B_{7L}(\pi_Z(X))$ can be extended to an $\mathcal{C}(X), z$ path where the addition also misses this set and the Hausdorff distance between $\pi_Z(X)$ and $\pi_X(Z)$ is L, we are done in this case.

Now suppose Y is the minimal element of $Y_K(X,Z)$. We apply [BBF, Lemma 3.9] to Y and in doing so deduce that every $\mathcal{C}(X), \mathcal{C}(Z)$ path meets $B_{7L}(\pi_Y(X))$. As $\operatorname{diam}(\pi_Y(X)) \leq L$ the result is complete. ☐

Fix a basepoint e lying in a piece $\mathcal{C}(E)$ and for each $Y \in \mathbf{Y}$, pick a closest point e_Y in $\mathcal{C}(Y)$ to e. Our construction will be dependent on some parameter R. Define $\mathbf{Y}^n := \big\{ Y \in \mathbf{Y} \;\big|\; d_{\mathcal{C}(\mathbf{Y})}(e, e_Y) \big\} \leq nR$ and set $\mathbf{Y}_n = \mathbf{Y}^n \setminus \mathbf{Y}^{n-1}$. For each $X \in \mathbf{Y}$ we define the *level* of X, $\operatorname{lev}(X)$, to be the unique n such that $X \in \mathbf{Y}_n$. Denote by $\mathcal{C}(\mathbf{Y}^n)$ the subgraph of $\mathcal{C}(\mathbf{Y})$ induced by all the pieces $\mathcal{C}(Y)$ with $Y \in \mathbf{Y}^n$ including all length L edges between them, so $\mathcal{C}(\mathbf{Y}^n) \supseteq B(nR; e)$ is connected. Given $X \in \mathbf{Y}_n$, we also fix a geodesic $\underline{g_X}$ from e_X to e in $\mathcal{C}(\mathbf{Y})$ and denote by c'_X the unique point on this geodesic at distance exactly $(n-1)R$ from e.

Lemma 2.2. *Given any $x \in \mathcal{C}(\mathbf{Y}^n)$, there is a $(1, 36L)$ quasi-geodesic from x to e contained entirely in $\mathcal{C}(\mathbf{Y}^n)$.*

Proof: $x \in \mathcal{C}(X)$ for some $X \in \mathbf{Y}^n$. Denote the concatenation of some geodesic from x to e_X with some geodesic from e_X to e by P. Note $P \subset \mathcal{C}(\mathbf{Y}^n)$ as pieces are totally geodesically embedded and $e_X \in B_{nR}(e)$ which is contained in $\mathcal{C}(\mathbf{Y}^n)$.

By lemma 2.1 any geodesic from x to e must pass within $18L$ of e_X. Hence

$$d_{\mathcal{C}(\mathbf{Y})}(x, e) \geq d_X(x, e_X) + d_{\mathcal{C}(\mathbf{Y})}(e_X, e) - 36L = |P| - 36L,$$

as $\mathcal{C}(X)$ is totally geodesically embedded. ☐

Lemma 2.3. *Let $X, Y \in \mathbf{Y}$ and suppose $d_{\mathcal{C}(\mathbf{Y})}(e_X, e) \geq d_{\mathcal{C}(\mathbf{Y})}(e_Y, e)$. Any path from $\mathcal{C}(X)$ to $\mathcal{C}(Y)$ passes within $72L$ of e_X.*

Proof: Suppose there is a path P from $x \in \mathcal{C}(X)$ to $y \in \mathcal{C}(Y)$ with $Y \in \mathbf{Y}^{n-1}$ which avoids the ball of radius $27L$ around e_X. If $d(e_X, e_Y) > 54L$ then any geodesic from e_Y to e avoids this ball, and as $\mathcal{C}(Y)$ is one-ended there is a path from y to e_Y also avoiding this ball, contradicting lemma 2.1.

Now consider a path of length at most $54L$ from e_X to e_Y, by lemma 2.1 some point on this path lies on a bottleneck for paths between $\mathcal{C}(X)$ and $\mathcal{C}(Y)$ and hence P contains a point within $72L$ of e_X. $\square$

3 Construction of the tree of metric spaces

We now introduce two equivalence relations on each level $\mathbf{Y}_n$, labelled $\sim_b$ and $\sim_n$. From here onwards we will assume that the space $\mathcal{C}(\mathbf{Y})$ is δ-hyperbolic. By this we mean given any geodesic triangle each edge lies in the δ neighbourhood of the union of the other two. We set $\delta(C)$ to be the corresponding constant obtained for any triangle with $(1, C)$–quasi-geodesic edges.

Given $X, Z \in \mathbf{Y}_n$ we write $X \sim_b Y$ if there exists a path from $\mathcal{C}(X)$ to $\mathcal{C}(Z)$ in $\mathcal{C}(\mathbf{Y})$ avoiding the ball of radius $C_0 = C_0(L, \delta)$ around e and write $X \sim_n Y$ if there exists a path from $\mathcal{C}(X)$ to $\mathcal{C}(Z)$ in $\mathcal{C}(\mathbf{Y})$ avoiding the C-tubular neighbourhood of $\mathcal{C}(\mathbf{Y}^{n-1})$. It is clear that $X \sim_b Z$ whenever $X \sim_n Z$. We denote equivalence classes by $[X]_b$ and $[X]_n$ respectively. The constant C_0 we choose is determined by the next lemma.

Lemma 3.1. *There exists some C_1 such that for all C_0 sufficiently large and all $Z \in [X]_n$, $d_{\mathcal{C}(\mathbf{Y})}(c'_X, c'_Z) \leq C_1$. Moreover, if $[X]_b \neq [X]_n$ for some $X \in \mathbf{Y}_n$, then there is a unique piece $Y \in \mathbf{Y}^{n-1}$ such that $d_{\mathcal{C}(\mathbf{Y})}(c'_Z, \mathcal{C}(Y)) \leq C_1$ for all $Z \in [X]_b$.*

Proof: Consider any standard path from $\mathcal{C}(X)$ to $\mathcal{C}(Z)$, notice that by lemma 2.1 the entry and exit points of the standard path into each piece lie within $18L$ of a point on $\underline{g_X} \cup \underline{g_Z}$, so must lie within a $36L$ neighbourhood of the restrictions of these geodesics to the range e_X to c'_X and e_Z to c'_Z respectively. We label these geodesics $\underline{g^c_X}$ and $\underline{g^c_Z}$ for brevity. There must be some $Y \in Y_K(X, Z)$ such that the standard path enters $\mathcal{C}(Z)$ close to $\underline{g^c_X}$ and exits close to $\underline{g^c_Z}$. Call the appropriate points on the two geodesics m_X and m_Z, these points must be at distance at least $C_0 - 18L$ away from c'_X and c'_Z respectively. As $\mathcal{C}(\mathbf{Y})$ is hyperbolic, $d_{\mathcal{C}(\mathbf{Y})}(m_X, m_Z) \geq d_{\mathcal{C}(\mathbf{Y})}(c'_X, c'_Z)$ whenever the second value is sufficiently large in terms of δ.

If $Y \notin \mathbf{Y}^{n-1}$, as is the case when $Z \in [X]_n$, we have a $(1, 144L)$ quasi-geodesic from m_X to m_Y obtained by travelling at most $36L$ to the entry point of the standard path into $\mathcal{C}(Z)$ then taking a geodesic in $\mathcal{C}(Z)$ to within $36L$ of m_Y and following a shortest path at the other end. This quasi-geodesic does not enter $B_{(n-1)R}(e)$.

Finally, we notice that if $d_{\mathcal{C}(\mathbf{Y})}(c'_X, c'_Z)$ is sufficiently large in terms of L then there will exist points on the quasi-geodesic edge which are not close to either of the geodesic edges $\underline{g_X}$ or $\underline{g_Y}$ contradicting the hyperbolicity of $\mathcal{C}(\mathbf{Y})$.

Alternatively, if $Y \in \mathbf{Y}^{n-1}$ then we define a quasi-geodesic $\underline{q_X}$ from e_X to e following $\underline{g_X}$ to m_X, moving to $\mathcal{C}(Y)$ using a path of length at most $36L$, then taking the quasi-geodesic via e_Y to e given by lemma 2.2. Note that this quasi-geodesic travels at least $C_0 - 54L$ steps closer to e while inside $\mathcal{C}(Y)$ so we obtain uniqueness of Y for fixed X and Z by lemma 2.1 when $C_0 > 72L$. As $\sim_b$ is an equivalence relation, this is sufficient to prove that such Y is unique for the entire equivalence class.

As the space is hyperbolic it is clear that for each $Z \in [X]_b$ the distance between c'_Z and any point on $\underline{q_Z}$ at distance $(n-1)R$ from e is small. By definition of $\underline{q_Z}$ such a point may be chosen which lies inside $\mathcal{C}(Y)$. ☐

Figure 1 below displays this argument. The uppermost path is guaranteed by the relation, the quasi-geodesics produced in the proof are represented by dotted lines.



Figure 1: The situations $X \sim_n Z$ and $X \sim_b Z$.

When such a Y exists we label this point on $\underline{q_Z}$ as c_Z. If $[X]_b = [X]_n$ then we must check one further thing: if some c'_Z (with $Z \in [X]_n$) lies within $\delta(36L)$ of some $\mathcal{C}(W)$ with $W \in \mathbf{Y}^{n-2}$ then we define c_Z to be some point in $\mathcal{C}(W)$ at distance at most $\delta(36L)$ from c'_Z. Otherwise, fix some X in this equivalence class and set $c_Z = c'_X$ for all $Z \in [X]_b$. In this way we define a point c_X for each $X \in \mathbf{Y} \setminus \{E\}$ at distance within a uniform constant of $(\mathrm{lev}(X) - 1)R$ from e. We denote by $c(X)$ the unique $W \in \mathbf{Y}^{n-1}$ such that $c_X \in \mathcal{C}(W)$.

With this done we are ready to construct a tree-graded space $\mathcal{T}(\mathbf{Y})$, which we do inductively with start point $\mathcal{T}(\mathbf{Y}^0) := \mathcal{C}(E)$.

To define $\mathcal{T}(\mathbf{Y}^n)$ we attach every piece $\mathcal{C}(X)$ with $X \in \mathbf{Y}_n$ to $\mathcal{T}(\mathbf{Y}^{n-1})$ as follows: add a single edge of length $d_{\mathcal{C}(\mathbf{Y})}(e_X, e) - d_{\mathcal{C}(\mathbf{Y})}(c_X, e)$ connecting $c_X \in \mathcal{T}(\mathbf{Y}^{n-1})$ to e_X. We define $\mathcal{T}(\mathbf{Y}) = \bigcup_{n \in \mathbb{N}} \mathcal{T}(\mathbf{Y}^n)$. This space is tree-graded in the sense of Druţu-Sapir, [DS05]. From this construction, we also define a simplicial tree $\mathcal{T}$ with vertex set $\mathbf{Y}$ and an edge between X and Y if and only if $c(X) = Y$ or $c(Y) = X$. We denote the standard metric on $\mathcal{T}$ by $d_{\mathcal{T}}$ and that on $\mathcal{T}(\mathbf{Y})$ by $d_{\mathcal{T}(\mathbf{Y})}$.

One key observation to make now is the following:

Lemma 3.2. *Suppose $X \sim_b Z$. Then $d_{\mathcal{T}}(X, Z) \leq 2$.* ☐

4 Results

First we prove the following.

Theorem 4.1. (Theorem 1) *The extension of identity maps on pieces defines a quasi-isometry $\phi : \mathcal{T}(\mathbf{Y}) \to \mathcal{C}(\mathbf{Y})$.*

The fact that $d_{\mathcal{C}(\mathbf{Y})}(\phi(x), \phi(y)) \leq K d_{\mathcal{T}(\mathbf{Y})}(x, y) + K$ follows immediately from the fact that geodesics in $\mathcal{T}(\mathbf{Y})$ define paths in $\mathcal{C}(\mathbf{Y})$ with at most a C_1 loss at each change of piece in $\mathcal{T}$.

We are left to check the other direction and this will be done by induction on $d_{\mathcal{T}}(X, Y)$.

Given $X, Y \in \mathbf{Y}$ we write the $\mathcal{T}$ geodesic from X to Y as

$$X = X_0, X_1, \ldots, X_k = Z = Y_l, Y_{l-1}, \ldots, Y_1, Y_0 = Y$$

where Z is the unique piece of minimal level.

Our base cases will be the situations $k = 0$ or $l = 0$ and $k = l = 1$. When $k = l = 0$ the result is obvious as pieces are totally geodesically embedded.

The other cases are dealt with by the next three lemmas.

Lemma 4.2. *Let $x \in \mathcal{C}(X)$ and suppose $y \in \mathcal{C}(Y)$ lies on some $\mathcal{T}(\mathbf{Y})$ geodesic from x to e. Then, providing R is sufficiently large,*

$$d_{\mathcal{T}(\mathbf{Y})}(x, y) \leq 2 d_{\mathcal{C}(\mathbf{Y})}(x, y) + C_2,$$

where the constant C_2 depends only on L and δ.

Proof: If $X = Y$ then $d_{\mathcal{T}(\mathbf{Y})}(x, y) = d_{\mathcal{C}(\mathbf{Y})}(x, y)$. We assume that the geodesic in $\mathcal{T}(\mathbf{Y})$ meets $n-1$ pieces (excluding X, Y) so $d_{\mathcal{C}(\mathbf{Y})}(x, y) \geq (n-2)R$. We write the pieces from X to Y in $\mathcal{T}$ as $X = X_0, X_1, \ldots, X_n = Y$.

For each $x \in \mathcal{C}(X)$ and suitable y we construct a path $\underline{q}(x, y)$ in $\mathcal{C}(\mathbf{Y})$ as follows: Consider any $\mathcal{C}(\mathbf{Y})$ geodesic $\underline{g_0}$ from x to e. By lemma 2.3 this passes within $72L$ of e_X, moreover, as $\mathcal{C}(\mathbf{Y})$ is hyperbolic and there is a geodesic from e_X to e passing through c'_X, $\underline{g_0}$ also contains a point m_0 within distance C'_2 of c'_X and hence within C''_2 of c_X. We follow $\underline{g_0}$ from x to m_0 then take any path P_0 of length at most C''_2 to c_X, then consider a $\mathcal{C}(\mathbf{Y})$ geodesic $\underline{g_1}$ from c_X to e. We repeat this process until we reach $c_{X_{n-1}}$ which lies in $\mathcal{C}(Y)$. By Lemma 2.2 there is a $(1, 18L)$ quasi-geodesic $\underline{q'}$ from $c_{X_{n-1}}$ to e passing through y, and we follow this path to y.

Then $\underline{q}(x, y)$ can be written as the concatenation

$$\underline{g_0}(x \to m_0), P_0, \underline{g_1}(c_{X_0} \to m_1), P_1, \ \ldots, \ \underline{g_n}(c_{X_{n-2}} \to m_{n-1}), P_{n-1}, \underline{q'}(c_{X_n} \to y).$$

Using the fact that pieces are totally geodesically embedded and assuming $R > 2C''_2$, we see that

$$\begin{aligned} d_{\mathcal{T}(\mathbf{Y})}(x, y) &\leq \left|\underline{q}(x, y)\right| \leq d_{\mathcal{C}(\mathbf{Y})}(x, y) + 2nC''_2 + 18L \\ &\leq 2 d_{\mathcal{C}(\mathbf{Y})}(x, y) + 6C''_2 + 18L. \end{aligned}$$

The result follows by setting $C_2 = 6C''_2 + 18L$. Moreover, by construction $\underline{q}(x, y)$ is a $(2, C_2)$ quasi-geodesic. ☐

Now we consider any two points $x \in \mathcal{C}(X)$, $y \in \mathcal{C}(Y)$. The following two examples demonstrate why we cannot expect this result to extend any further without introducing an additive error dependent on R. The first picture dictates its necessity in lemma 4.3 while the second demonstrates the need for these relations and explains the additive error in lemma 4.4.



Figure 2: The necessity of R-dependence in the additive error

To match the hypotheses we could assume all lines in the right hand picture of figure 2 represent one-ended graphs and that each pair of black dots joined by an edge containing no other black dot has length 1. This space is then quasi-isometric to a tree-graded space.

Lemma 4.3. *Suppose in the situation above* $l = 0$*, so* $Y = Z$*. There exists some constant* $C_3 = C_3(L, \delta)$ *such that*

$$d_{\mathcal{T}(\mathbf{Y})}(x, y) \leq d_{\mathcal{C}(\mathbf{Y})}(x, y) + d_{\mathcal{T}}(X, Y)C_3 + 2R.$$

Proof: We proceed by induction on k. If $k = 1$ the result follows from lemma 2.3. Now suppose $k > 1$. By the inductive hypothesis

$$d_{\mathcal{T}(\mathbf{Y})}(c_X, y) \leq d_{\mathcal{C}(\mathbf{Y})}(c_X, y) + (d_{\mathcal{T}}(X, Y) - 1)C_3 + 2R.$$

Suppose some geodesic from x to y meets $B_{\delta(36L)}(c_X)$. Then $d_{\mathcal{C}(\mathbf{Y})}(x, y) \geq d_{\mathcal{C}(\mathbf{Y})}(x, c_X) + d_{\mathcal{C}(\mathbf{Y})}(c_X, y) - 4\delta(36L)$.

Combining these facts we see that

$$\begin{aligned} d_{\mathcal{T}(\mathbf{Y})}(x, y) &\leq d_{\mathcal{T}(\mathbf{Y})}(x, c_X) + d_{\mathcal{C}(\mathbf{Y})}(c_X, y) + (d_{\mathcal{T}}(X, Y) - 1)C_3 + 2R \\ &\leq d_{\mathcal{C}(\mathbf{Y})}(x, c_X) + (2C_1 + 36L) + \\ &\quad\; d_{\mathcal{C}(\mathbf{Y})}(c_X, y) + (d_{\mathcal{T}}(X, Y) - 1)C_3 + 2R \\ &\leq d_{\mathcal{C}(\mathbf{Y})}(x, y) + (d_{\mathcal{T}}(X, Y) - 1)C_3 + 2R + \\ &\quad\; 2C_1 + 36L + 2\delta(36L). \end{aligned}$$

So the result holds providing $C_3 \geq 2C_1 + 36L + 2\delta(36L)$.

Alternatively, if all geodesics from x to y in $\mathcal{C}(\mathbf{Y})$ avoid $B_{\delta(36L)}(c_X)$, then (considering the $(1, 36L)$ quasi-geodesics from x and y to e) by hyperbolicity c'_X must lie within $\delta(36L)$ of the geodesic y to e_Y, as $\text{lev}(Y) \leq \text{lev}(X) - 2$. Therefore, c'_X lies within $\delta(36L)$ of $\mathcal{C}(Y)$ with $Y \in \mathbf{Y}^{n-2}$ so by construction $c_X \in \mathcal{C}(Y)$. Hence, $d_{\mathcal{T}}(X, Y) = 1$ which is a contradiction. ☐

The case $k = l = 1$ is dealt with next.

Lemma 4.4. *Suppose $k = l = 1$. There exists some constant $C_4 = C_4(L, \delta)$ such that*

$$d_{\mathcal{T}(\mathbf{Y})}(x, y) \leq d_{\mathcal{C}(\mathbf{Y})}(x, y) + 4R + C_4.$$

Proof: We assume that $d(e, e_X) \geq d(e, e_Y)$ so any path from x to y must meet $B(e_X; 72L)$ by lemma 2.3. If some geodesic from x to y meets the ball around c_X of radius $\delta(72L + 2C''_2)$ then we are done by lemma 4.3 for sufficiently large C_4. Otherwise, consider a $(1, 72L + 2C''_2)$ quasi-geodesic triangle comprising the paths $\underline{q}(x, e_Z)$, $\underline{q}(y, e_Z)$ and some geodesic from x to y. Then $d_{\mathcal{C}(\mathbf{Y})}(e_Y, e) \geq d_{\mathcal{C}(\mathbf{Y})}(c_X, e) - (18L + 2C''_2)$ as the geodesics c_X to e_Z in $\mathcal{C}(Z)$ and y to e_Y in $\mathcal{C}(Y)$ cannot remain close for longer than $18L$ without contradicting lemma 2.1. This implies $\text{lev}(X) \geq \text{lev}(Y) \geq \text{lev}(X) - 1$ for R sufficiently large. Therefore, we notice that $d_Z(c_X, c_Y) \leq R + C'_4$ for some uniform constant C'_4, so using this quasi-geodesic triangle we see that

$$d_{\mathcal{T}(\mathbf{Y})}(x, y) \leq d_{\mathcal{C}(\mathbf{Y})}(x, y) + 4R + 2C'_4 + 2\delta(72L + 2C''_2).$$

We complete the proof by ensuring $C_4 \geq 2C'_4 + 2\delta(72L + 2C''_2)$. ☐

Now we are ready to complete the proof of Theorem 4.1.

We proceed by induction on $d_{\mathcal{T}}(X, Y)$ and note that our base cases ensure we can assume $d_{\mathcal{T}}(X, Y) \geq 3$ and $k, l \geq 1$, under the hypothesis

$$d_{\mathcal{T}(\mathbf{Y})}(x, y) \leq d_{\mathcal{C}(\mathbf{Y})}(x, y) + 4R + d_{\mathcal{T}}(X, Y)C_5.$$

We prove that this equation holds for some uniform $C_5 = C_5(L, \delta)$.

In the first instance assume that $d_{\mathcal{C}(\mathbf{Y})}(\{c_X, c_Y\}, \underline{g}) \geq C'_5$ for some geodesic $\underline{g}$ between e_X and e_Y and some $C'_5 = C'_5(L, \delta)$ which is sufficiently large.

Then considering the quasi-geodesic triangle compromising $\underline{g}$ and the two quasi-geodesics defined in lemma 4.2 from e_X and e_Y to e respectively, we deduce that as $\mathcal{C}(\mathbf{Y})$ is hyperbolic, c_X and c_Y are uniformly close in $\mathcal{C}(\mathbf{Y})$, so $\text{lev}(X) = \text{lev}(Y)$ providing R is sufficiently large. Moreover, $\underline{g}$ defines a path from $\mathcal{C}(X)$ to $\mathcal{C}(Y)$ lying outside the ball of radius $(n-1)R + C_1$ around e, thus $X \sim_b Y$. But this is a contradiction as $d_{\mathcal{T}}(X, Y) \leq 2$ by lemma 3.2.

So, we are left with the case where $d_{\mathcal{C}(\mathbf{Y})}(\{c_X, c_Y\}, \underline{g}) \leq C'_5$, assume without loss of generality that c_X lies close to $\underline{g}$. Then there are uniform $(1, C''_5)$ quasi-geodesics from $x \in \mathcal{C}(X)$ to $y \in \mathcal{C}(Y)$ obtained by travelling from x to e_X geodesically then following $\underline{g}$ to some point close to c_X (guaranteed by assumption), moving to c_X and then following any geodesic from c_X to y. The inductive hypothesis gives $d_{\mathcal{T}(\mathbf{Y})}(c_X, y) \leq d_{\mathcal{C}(\mathbf{Y})}(c_X, y) + 4R + (d_{\mathcal{T}}(X, Y) - 1)C_5$, so following this quasi-geodesic we see that

$$d_{\mathcal{T}(\mathbf{Y})}(x, y) \leq d_{\mathcal{C}(\mathbf{Y})}(x, y) + 4R + d_{\mathcal{T}}(X, Y)C_5$$

providing $C_5 \geq 2C''_5$. As this constant C_5 does not depend on $d_{\mathcal{T}}(X, Y)$ this completes the induction.

Then taking $R \geq 2C_5$ and noticing that $d_{\mathcal{T}(\mathbf{Y})}(x,y) \geq R(d_{\mathcal{T}}(X,Y) - 4)$ we have

$$d_{\mathcal{T}(\mathbf{Y})}(x,y) \leq 2d_{\mathcal{C}(\mathbf{Y})}(x,y) + 8R + 16C,$$

where $C = \max\{C_3, C_4, C_5\}$ so ϕ is a quasi-isometry. □

We then obtain

Theorem 4.5. (Theorem 2) *The mapping class group of any compact surface has finite Assouad-Nagata dimension.*

Proof: Let $S = S_{g,n}$ be a compact surface. If $3g + n - 4 \leq 0$ then $MCG(S)$ is hyperbolic and the result follows. We now assume $3g + n - 4 > 0$. By [BBF], such mapping class groups quasi-isometrically embed into a finite product of spaces $\mathcal{C}(\mathbf{Y})$ where all the curve complexes $\mathcal{C}(Y)$ in any $\mathcal{C}(\mathbf{Y})$ are isometric. If one of the complexes $\mathcal{C}(Y)$ is not one-ended, then the transitive action of a mapping class group of some subsurface of S on its 0-skeleton forces each $\mathcal{C}(Y)$ to be a quasi-tree, (it is simple to check such a space has Manning's bottleneck property [Man05]) so $\mathcal{C}(\mathbf{Y})$ is also a quasi-tree by [BBF, Theorem 3.10]. In such situations we define $\mathcal{T}(\mathbf{Y})$ to be some appropriate tree which is quasi-isometric to $\mathcal{T}(\mathbf{Y})$. Then we deduce from theorem 1 that mapping class groups quasi-isometrically embed into finite products of trees and tree-graded spaces in which all pieces have Bowditch's tight geodesic property [Bow08]. Thus all the $\mathcal{T}(\mathbf{Y})$ have finite asymptotic Assouad-Nagata dimension [BF08, BH], so we deduce that the mapping class group does as well [LS05]. Finally, for all discrete metric spaces the asymptotic Assouad-Nagata dimension equals the Assouad-Nagata dimension, so the result follows. □

As mentioned previously, this implies that mapping class groups have ℓ^p compression exponent 1 for all $p \in [1, \infty)$, however, using additional tools we can provide explicit embeddings of mapping class groups. The first requirement is a method of embedding curve complexes into ℓ^p spaces, which requires the Masur-Minsky theorem that such spaces are δ-hyperbolic and Bowditch's results on finiteness properties of tight geodesics.

Theorem 4.6. *Let X be a hyperbolic simplicial graph satisfying Bowditch's tight geodesic property. Then $\alpha_p^*(X) = 1$.*

We actually prove the stronger result that whenever $f : \mathbb{N} \to \mathbb{N}$ is a function satisfying the properties

- (concavity) $f(n+1) - f(n) \leq f(n) - f(n-1)$ for all $n \geq 1$,
- (C_p) $\sum_n \frac{1}{n}\left(\frac{f(n)}{n}\right)^p < \infty$,

then there is an embedding ϕ of X into an ℓ^p space with

$$f(d(x,y)) \preceq \|\phi(x) - \phi(y)\|_p \preceq d(x,y).$$

From this result we can deduce the same conclusion for all curve graphs of surfaces in the following manner. When the complexity $\kappa(S) = 3g + n - 4 > 0$, the curve graph is quasi-isometric to the 1-skeleton of the standard curve complex $\mathcal{C}(S)$, so Bowditch's result on tight geodesics applies and we can use theorem 4.6. When the complexity is lower than this all curve complexes are quasi-trees, so admit a suitable embedding via [Gal08, Tes11] and an explicit embedding via [Man05, Hum11].

The following lemma will be required.

Lemma 4.7. *Let X be the 0-skeleton of a δ-hyperbolic graph and let $e \in X$. Then there exists some constant $K = K(\delta)$, such that for all $n \geq 3\delta$, for all $x, y \in X$ with $d(x,e) \geq n$ and $d(x,y) \leq \frac{n}{4}$, for all geodesics $\underline{g_0}$ from x to e, $\underline{g}$ from y to e and $p \in \underline{g}([n, 2n])$,*

$$d\left(p, \underline{g_0}\left(\left[\frac{n}{2}, \frac{5n}{2}\right]\right)\right) \leq K.$$

See [Hum11, Lemma 2.1]. □

In what follows we only deal with geodesics which are tight in the sense of Bowditch [Bow08]. We recall here Bowditch's notation. $\mathcal{L}_T(a,b)$ denotes the set of all tight geodesics connecting a and b, of which there are finitely many (and at least one [MM00]). Moreover, we set

$$\mathcal{L}_T(A,b) = \bigcup_{a \in A} \mathcal{L}_T(a,b).$$

Fix a basepoint $e \in \mathcal{C}(S)$, we define the set of restricted geodesics

$$\underline{G_{x,k,n}} = \bigcup_{\underline{g} \in \mathcal{L}_T\left(\overline{B_k(x)}, e\right)} \left\{\underline{g}([n, 2n])\right\}.$$

We define $F_{x,k,n}$ to be the set of all points in $\mathcal{C}(X)$ lying on some $\underline{g} \in \underline{G_{x,k,n}}$ but not in $B_{3\delta}(e)$ and set $F(x,k,n)$ to be the characteristic function of $F_{x,k,n}$.

Lemma 4.8. *There exists some n_0 such that for all $n \geq n_0$, $F_{x,k,n}$ is a finite set, so $F(x,k,n) \in \ell^p(\mathcal{C}(S))$.*

Proof: This follows from [Bow08, Theorem 1.1] and lemma 4.7. □

Next define

$$H(x,n) = \frac{1}{n} \sum_{k \leq \frac{n}{4}} F(x,k,n).$$

The following lemma captures the necessary properties of these functions.

Lemma 4.9. *There exists some constant C such that for all $x, y \in \mathcal{C}(S)$, $k \leq \frac{n}{4}$ and $n \geq n_0$,*

(i) $\|F(x,k,n)\|_p^p \leq Cn$, *and if* $d(x,e) \geq 2n$, *then* $\|F(x,k,n)\|_p^p \geq n - 3\delta$,

(ii) $\|H(x,k,n)\|_p^p \leq Cn$, *and if* $d(x,e) \geq 2n$, *then* $\|H(x,k,n)\|_p^p \geq \frac{n-3\delta}{4}$,

(iii) *if* $d(x,y) \leq R$, *then* $\|H(x,n) - H(y,n)\|_p \leq 2C(R+1)n^{-\frac{p-1}{p}}$.

The proofs are so similar to those of [Hum11, Lemmas 2.3-2.5] that we omit additional details here. □

We can now define our embedding $\phi : \mathcal{C}(S) \to \bigoplus_n \ell^p(\mathcal{C}(S))$, by setting

$$\phi(x) = \sum_{n \geq \log_2(n_0+3)} \frac{f(2^n)}{2^{\frac{n}{p}}} H(x, 2^n).$$

The result then follows in exactly the same way as [Hum11]. $\square$

With this we can prove the final theorem of the paper.

Theorem 4.10. *For each $p \in [1, \infty)$ there is an explicit embedding of each mapping class group into some ℓ^p space exhibiting compression exponent 1.*

Proof: Given the embeddings of curve complexes given by theorem 4.6 we use the tree-graded space embedding from [Hum11] to deduce that the spaces $\mathcal{T}(\mathbf{Y})$ have compression exponent 1 (for the exact result we need an additional constraint on functions f in theorem 4.6 explained in [Hum11]). Combining these results we obtain an explicit embedding

$$MCG(S) \to \prod \mathcal{C}(\mathbf{Y}) \to \prod \mathcal{T}(\mathbf{Y}) \to \ell^p(\mathbb{N}),$$

exhibiting compression exponent 1.

References

[Ass82] Patrice Assouad. Sur la distance de Nagata. *C. R. Acad. Sci. Paris Sér. I Math.*, 294(1):31–34, 1982.

[BBF] Mladen Bestvina, Ken Bromberg, and Koji Fujiwara. The asymptotic dimension of mapping class groups is finite. (preprint). Available from arXiv:1006.1939v2.

[BDHM09] Nikolay Brodskiy, Jerzy Dydak, Jose Higes, and Atish Mitra. Assouad-Nagata dimension via Lipschitz extensions. *Israel J. Math.*, 171:405–423, 2009.

[BDL] Nikolay Brodskiy, Jerzy Dydak, and Urs Lang. Assouad-nagata dimension of wreath products of groups. Preprint available from http://arxiv.org/abs/math/0611331.

[BDLM08] Nikolay Brodskiy, Jerzy Dydak, Michael Levin, and Atish Mitra. A Hurewicz theorem for the Assouad-Nagata dimension. *J. Lond. Math. Soc. (2)*, 77(3):741–756, 2008.

[Beh04] Jason A. Behrstock. *Asymptotic Geometry of the Mapping Class Group and Teichmuller Space.* PhD thesis, SUNY Stony Brook, 2004.

[BF08] Gregory C. Bell and Koji Fujiwara. The asymptotic dimension of a curve graph is finite. *J. Lond. Math. Soc. (2)*, 77(1):33–50, 2008.

[BH] Nikolay Brodskiy and Jose Higes. Assouad-Nagata dimension of tree-graded spaces. arXiv:0910.2378v1.

[BM11] Jason A. Behrstock and Yair N. Minsky. Centroids and the rapid decay property in mapping class groups. *J. Lond. Math. Soc. (2)*, 84(3):765–784, 2011.

[Bow08] Brian H. Bowditch. Tight geodesics in the curve complex. *Invent. Math.*, 171(2):281–300, 2008.

[DH08] Jerzy Dydak and Jose Higes. Asymptotic cones and Assouad-Nagata dimension. *Proc. Amer. Math. Soc.*, 136(6):2225–2233, 2008.

[DS05] Cornelia Druţu and Mark Sapir. Tree-graded spaces and asymptotic cones of groups. *Topology*, 44(5):959–1058, 2005. With an appendix by Denis Osin and Mark Sapir.

[Gal08] Światosław R. Gal. Asymptotic dimension and uniform embeddings. *Groups Geom. Dyn.*, 2(1):63–84, 2008.

[Gro93] Misha Gromov. Asymptotic invariants of infinite groups. In *Geometric group theory, Vol. 2 (Sussex, 1991)*, volume 182 of *London Math. Soc. Lecture Note Ser.*, pages 1–295. Cambridge Univ. Press, Cambridge, 1993.

[Ham09] Ursula Hamenstädt. Geometry of the mapping class groups. I. Boundary amenability. *Invent. Math.*, 175(3):545–609, 2009.

[HR00] Nigel Higson and John Roe. Amenable group actions and the Novikov conjecture. *J. Reine Angew. Math.*, 519:143–153, 2000.

[Hum11] David Hume. Direct embeddings of relatively hyperbolic groups with optimal ℓ^p compression exponent. Preprint (arXiv:1111.6013v1), 2011.

[Kid08] Yoshikata Kida. The mapping class group from the viewpoint of measure equivalence theory. *Mem. Amer. Math. Soc.*, 196(916):viii+190, 2008.

[LS05] Urs Lang and Thilo Schlichenmaier. Nagata dimension, quasisymmetric embeddings, and Lipschitz extensions. *Int. Math. Res. Not.*, (58):3625–3655, 2005.

[Man05] Jason Fox Manning. Geometry of pseudocharacters. *Geom. Topol.*, 9:1147–1185 (electronic), 2005.

[MM99] Howard A. Masur and Yair N. Minsky. Geometry of the complex of curves. I. Hyperbolicity. *Invent. Math.*, 138(1):103–149, 1999.

[MM00] Howard A. Masur and Yair N. Minsky. Geometry of the complex of curves. II. Hierarchical structure. *Geom. Funct. Anal.*, 10(4):902–974, 2000.

[Now07] Piotr W. Nowak. On exactness and isoperimetric profiles of discrete groups. *J. Funct. Anal.*, 243(1):323–344, 2007.

[NP11] Asaf Naor and Yuval Peres. L_p compression, traveling salesmen, and stable walks. *Duke Mathematical Journal*, vol. 157 no.1:pages 53–108, 2011.

[Sch05] Thilo Dirk Schlichenmaier. *A quasisymmetrically invariant notion of dimension and absolute Lipschitz retracts*. ProQuest LLC, Ann Arbor, MI, 2005. Thesis (Dr.sc.math.)–Eidgenoessische Technische Hochschule Zuerich (Switzerland).

[Sch11] Saul Schleimer. The end of the curve complex. *Groups, Geometry, and Dynamics*, no.1:169–176, 2011.

[Tes11] Romain Tessera. Asymptotic isoperimetry on groups and uniform embeddings into Banach spaces. *Comment. Math. Helv.*, 86(3):499–535, 2011.

[Yu00] Guoliang Yu. The coarse Baum-Connes conjecture for spaces which admit a uniform embedding into Hilbert space. *Invent. Math.*, 139(1):201–240, 2000.